

## Comparison the Effects of Listening to Three Types of Music during Exercise on Heart Rate, Blood Pressure, Rating of Perceived Exertion and Fatigue Onset Time

Mohammad Reza RamezanPour<sup>1\*</sup>, Amir Moghaddam<sup>1</sup>, Elaheh Sadifar<sup>2</sup>

<sup>1</sup>Department of Physical Education, Mashhad Branch, Islamic Azad University, Mashhad, Iran

<sup>2</sup>MSc in Physical Education and Sport Sciences

Received 4 September 2011

Accepted 15 December 2011

### Abstract

**Introduction:** The present study aimed at investigating the effects of listening to three types of music during exercise on heart rate, blood pressure, rating of perceived exertion (RPE) and fatigue onset time.

**Material and Methods:** 15 female students of physical education with the mean age of  $21.93 \pm 2.43$  year's height of  $161.8 \pm 5.22$  centimeters and weight of  $54.46 \pm 7.27$  kilograms were randomly selected as the participants from 45 volunteers. After warm-up, the participants started pedaling at 80-85% of maximum heart rate until exhaustion. The participants performed this exercise on four different days (with a one-week interval). The first exercise session was performed without listening to music, the second session was accompanied with some exciting music, the third session was held with soft, relaxing music and the last one with arbitrary music. RPE was recorded every 3 minutes during the exercise. After exhaustion, RPE, heart rate, systolic and diastolic blood pressure, and fatigue onset time were recorded. One-way ANOVA (F) was used to analyze the data and LSD test was run to compare the differences ( $P \leq 0.05$ ).

**Results:** Listening to exciting music during exercise significantly affects heart rate, RPE and fatigue onset time but not systolic and diastolic blood pressure. Listening to relaxing music during exercise significantly affects heart rate and systolic blood pressure but not diastolic blood pressure, RPE and fatigue onset time. Listening to arbitrary music, as selected by the participants themselves, during exercise significantly affects heart rate, systolic blood pressure, RPE and fatigue onset time but not diastolic blood pressure.

**Discussion and Conclusion:** Understanding the effects of music genre on the variables examined may help the athletes select a proper genre of music based on the type of exercise they do. For example, listening to exciting and arbitrary types of music significantly increases heart rate while relaxing music significantly decreases heart rate.

**Keywords:** Music, Heart rate, Blood pressure, Rating of perceived exertion, Fatigue onset time

### Introduction

Art has always served human's perfectionist and charm-seeking needs. As a soothing element, it has also helped humans tolerate their pains and sufferings. Music is a type of art which conveys human emotions, feelings and perception nonverbally. Music is often regarded as an original source of provocation and inspiration in sports and athletes' exercises. There have yet been fundamental discussions about music and its relation to other arts and sports phenomena. According to some scholars, exciting music increases agitation while soft music often blocks the transmission of negative information to the nervous system [1, 2, and 3]. Based on previous studies, different genres of music may affect the athletes' performance while the results have sometimes been contradictory and sometimes confirmatory [3]. Besides, music may encourage the athletes to continue

their exercise [4] and delay fatigue via reducing rate of perceived exertion on the body [5-7].

Based on the present theories, music narrows down the attention and distracts the individual from activity-induced fatigue, removes mental agitation, serves as an exciting or soothing means before or during the activity, and helps the body respond to the rhythmic components of music in sub maximal activities [8].

Copland and Franks (1991) investigated the effects of different genres of music on heart rate, RPE and fatigue time in 24 college students while they were running on the treadmill. They reported that exciting music increases heart rate to a higher extent. Moreover, fatigue time during exercise with exciting music was delayed compared to the control group [9]. Branda et al (1991) investigated the effects of exciting music and soft music types on the heart rate of athletes running on the treadmill until exhaustion. The athletes' heart rate was recorded every 30 seconds. The results showed that soft music, as compared to exciting music and

---

\* Corresponding author E-mail: ramezanpour@mshdiau.ac.ir

control group (no music), decrease heart rate at the 1<sup>st</sup> and 6<sup>th</sup> minutes but increased the heart rate during the exercise and immediately before exhaustion. The athletes' heart rate was higher in the control group and lower than exciting music conditions comparing to soft music [10]. Khorram (2005) studied the effects of exciting and classical music on salivary cortisol concentration and RPE during a session of training to the exhaustion stage on the treadmill at the 70% of maximum heart rate. The results showed that listening to music significantly increased the running time in the athletes but salivary cortisol concentration and RPE were significantly lower in them comparing to the control group. There was no significant relationship between salivary cortisol concentration and RPE after listening to music [5]. Fathollahi (2006) investigated the effects of two types of music on the running time, salivary cortisol and RPE in a session of training in active women. The results showed that listening to the exciting and soft music did not significantly affect the running time, cortisol concentration and RPE. As well, there was no significant relationship between salivary cortisol concentration and RPE after listening to soft music; however, this relationship was significant after listening to the exciting music [11]. Yamashita et al (2006) studied the effects of listening to music during exercise on RPE, heart rate and automatic nervous system. The purpose of their study was to investigate the effects of music on RPE and automatic nervous system both before and after sub maximal activity. The results revealed that, at 60% of VO<sub>2</sub> max, there was no significant difference in the RPE between the activities accompanied with music, and the activities without music. However, at the 40% of VO<sub>2</sub> max, RPE was lower in the activities accompanied with music compared to activities with no music. The heart rate and blood pressure at 40% and 60% of VO<sub>2</sub> max were significantly different between with music and without music conditions. This implies that music may cause agitation at low intensity exercise but it may not affect the automatic nervous system [12].

Due to the variety of music genres and their different effects on physiological and psychological factors and regarding the individuals' age, gender and mood as well as the time of listening, playback

devices, temperament, customs, traditions, musical and cultural characteristics, personal preferences, the intensity of physical activities and different patterns of training programs, more investigations on this issue are needed. In this regard, the researchers posed the following research questions to be investigated in this study:

1. Does listening to music during exercise affect the physiological factors such as heart rate, blood pressure, RPE and fatigue onset time in the female students of physical education?

2. Do different types of music exert the same influence on the above-mentioned physiological factors?

### Material and Methods

This is a practical study which was conducted with three experimental (exciting, relaxing and arbitrary music) and one control mode. The population of the study consisted of **150** female students of physical education in Bojnord Islamic Azad University.

**Participants:** From the population, **15** female students with the mean age of  $21.93 \pm 2.43$  years, mean height of  $161.8 \pm 5.22$  centimeters and mean weight of  $54.46 \pm 7.27$  kilograms were randomly selected as the participants. They were all living in the university dorm and took regular exercises 3 days a week on average. No one had a history of mental, ontological and hormonal disorders.

**Training program:** The training protocol consisted of pedaling the exercise bike at the intensity of 80-85% of maximum heart rate till exhaustion. In this regard, the participants exercised on the exercise bike on four different days (with a one-week interval) The first exercise session was performed without listening to music, the second session was accompanied with some exciting music, the third session was held with soft, relaxing music, and the last session with arbitrary music. All the participants listened to music over headphones. The main exercise protocol started at 20 Watts and the pressure was increased by 20 Watts every 3 minutes. The participants' pedaling rhythm was 80 rpm (Table 1).

**Table1.** Participants' training program in each exercise session

music type	Pedal Load (Watt)	Velocity (Round / min)	Increase Work load until Exhaustion (watt)
Exciting	20	80	20
Relaxing	20	80	20
Arbitrary	20	80	20
Control	20	80	20

**Data collection and procedure:** The participants were informed about the research process in a meeting. Then they were asked to fill out a form requiring their personal information. Next, their blood pressure and pulse were measured at rest. Then, having done a 10-minute warm-up (1 minute of fast walk, 1 minute of jogging, 8 minutes of static and dynamic stretching exercises), they did the pedaling test. A digital wrist barometer was used to measure heart rate and blood pressure. Systolic and diastolic blood pressure and heart rate were recorded at rest and immediately after the test; the participants' RPE was measured using the Borg 20-point scale. To this end, after the warm-up and at the earlier stages of the test while they were pedaling, the participants were asked to announce a point on the Borg 20-point scale which best described their perceived exertion. Their perceived exertion was recorded every 3 minutes from the start of the test to the exhaustion. Fatigue onset time was determined based on the disruption of the pedaling rhythm. The initial pedaling rhythm was 80 rpm. Once this rhythm was disrupted and the participant could not restore it, it was regarded as the fatigue onset time for the participant. At the end of every 3 minutes, before the increase of exertion, RPE was recorded on the Borg scale. At the time of exhaustion, RPE, heart rate, systolic and diastolic blood pressure and fatigue onset time were recorded again.

**Instrument:** The instruments of the study included an informed consent to be filled with

personal information, JEMIS digital chronometer made in Taiwan, UNIVERSAL full automatic exercise bike made in Taiwan, OMRON digital wrist barometer and pulse meter Model RX-3 made in China, MP3 player and headphones and some pieces of exciting and relaxing music.

**Data analysis:** Descriptive statistics including mean and standard deviation was used to categorize the data and to draw diagrams. One-way ANOVA (F) was run to analyze the data and LSD test was run to examine the differences. The level of significance was set at  $P \leq 0.05$  for all the tests.

## Results

As it is shown in Table 2, there are significant differences between variables of heart rate ( $p=0.001$ ), systolic blood pressure ( $p=0.003$ ), rating of perceived exertion ( $p=0.001$ ) and fatigue onset time ( $p=0.001$ ) in different conditions (listening to music and without music).

The results of Post hoc test (LSD) are delineated as follows:

1. LSD for the heart rate showed that the differences in the heart rate mean in all the three conditions with music were significant except for the difference between exciting and arbitrary music ( $P \leq 0.05$ ).

2. LSD for the Systolic Pressure showed that differences in the mean Systolic Pressure in all the three conditions with music were significant except for the difference between the control group and the exciting and arbitrary music and between the arbitrary music and the control group ( $P \leq 0.05$ ).

**Table 2:** Mean, comparative F and LSD test among listening to three types of music during exercise

variables	conditions	Mean $\pm$ SD	F	P-Value	Post hoc test (LSD)
<b>Heart rate (b/ min)</b>	Exciting music	155.53 $\pm$ 7.47	9.028	0.001	Significant in all conditions except between exciting with arbitrary music
	Relaxing music	148.13 $\pm$ 8.39			
	Arbitrary music	154.93 $\pm$ 8.69			
	control	152.33 $\pm$ 6.98			
<b>Systolic Pressure (mmHg)</b>	Exciting music	13.86 $\pm$ .875	6.410	0.003	Sig. in all conditions except between exciting with music arbitrary and both with control group
	Relaxing music	12.96 $\pm$ .676			
	Arbitrary music	13.57 $\pm$ .537			
	control	13.363 $\pm$ .486			
<b>Diastolic Pressure (mmHg)</b>	Exciting music	8.080 $\pm$ 1.068	0.312	0.869	Not sig.
	Relaxing music	8.226 $\pm$ 0.658			
	Arbitrary music	8.240 $\pm$ .572			
	control	8.273 $\pm$ 0.756			
<b>perceived exertion rate (score)</b>	Exciting music	13.53 $\pm$ 1.302	10.098	0.001	Sig. in all conditions except between arbitrary music with control group
	Relaxing music	14.80 $\pm$ 1.264			
	Arbitrary music	13.86 $\pm$ 1.187			
	control	14.66 $\pm$ 1.170			
<b>fatigue onset time (min)</b>	Exciting music	8.18 $\pm$ 2.050	8.164	0.001	Sig. in all conditions except between exciting with arbitrary and relaxing music with control group
	Relaxing music	7.21 $\pm$ 1.400			
	Arbitrary music	7.59 $\pm$ 1.590			
	control	7.19 $\pm$ 1.190			

3. Also LSD for the rating of perceived exertion showed that the differences in the mean rating of perceived exertion in the three conditions accompanied with music were significant except for the difference between arbitrary music and the control group ( $P \leq 0.05$ ).

4. And LSD for fatigue onset time showed that the differences in the mean fatigue onset time in all the conditions accompanied with music were significant, except for the difference between the exciting and arbitrary music, and between relaxing music and the control group ( $P \leq 0.05$ ).

### Discussion and Conclusion

The results showed that music affects the heart rate of female students of physical education during exercise. Exciting and arbitrary music significantly increase heart rate while relaxing music significantly decreases heart rate. This finding is consistent with the findings of Mohammed Zadeh and Ahmedi [13], Torabi [14], Liptalk [15], Davis [16], Branda et al [10], Copland and Franks [9], Brownley et al [17], Knight and Richard [18], Nethery [7] and Yamashita et al [12]. However, it is inconsistent with the findings of Nikbakhsh [9], Szabo et al [20], Lee Crust [21] and Yamamutu et al [22].

The effects of relaxing music on heart rate reduction may be attributed to the increase in plasma norepinephrine [9], decrease in muscular tension [23], decrease in cardiac pressure [24], decrease in factors associated with pain and discomfort [9] and/or the nervous effects of relaxing music which are not yet fully known. On the other hand, the effects of exciting music on heart rate increase may relate to the increase in blood norepinephrine [9], increase in cortisol concentration [25], decrease of muscular tension [23] and the nervous effects of exciting music. Besides, short movements of different limbs coordinating with music rhythm may increase these factors as well [2].

The results showed that listening to music during exercise affects the systolic but not diastolic blood pressure in the female students of physical education. Although exciting music increased the mean systolic blood pressure, the increase was not significant. However, relaxing and arbitrary music significantly affected systolic blood pressure. This is consistent with the findings of Mohammed Zadeh and Ahmedi [13], Davis [16], Knight and Richard [18] and Yamashita et al [12].

Listening to music during exercise affects RPE

in the female students of physical education. Accordingly, exciting and arbitrary music decreased RPE, but exercising with relaxing music or without music increased the RPE in the students. This is consistent with the findings of Khoram [5], Fathollahi [11], Ghaderi [26], Ahmedi [27], Szmedra et al [3], Knight and Richard [18], Meeks et al [28], Nethery [7], Maraki et al [29] and Yamashita et al [12], but it is inconsistent with findings of Nikbakhsh [19] and Lee Crust [21].

According to Rejeski a combination of psychological and physiological factors affect RPE during parallel processing of information. In this regard, sensory information and stimuli are subconsciously processed in parallel [30]. Thus, sensory information like exertion or stimuli like encouragement and workload-induced worries affect the RPE in the individual. Rejeski suggested that physiological factors are significantly more effective than psychological responses in heavy exercises. Besides, external stimuli like music may be very effective in low intensity exercise [30]. According to Boutcher and Trenske, listening to music during exercise may block the unfavorable feedback from peripheral and central factors to the central nervous system allowing the individual to feel more comfortable during the exercise [6].

The present findings revealed that listening to music during exercise affects fatigue onset time in the female students of physical education. Both exciting and arbitrary music delayed the fatigue onset time, that is, these types of music increased the pedaling time. However, listening to either relaxing or no music decreased the fatigue onset time and consequently reduced the pedaling time. This is consistent with the findings of Khoram [5], Ghaderi [26], Torabi [14], Liptalk [15], Davis [16], Meeks et al [28], Copland and Franks [9] and Szabo et al [20], but is inconsistent with the findings of Fathollahi [11] and Yamamutu et al [22].

Research has shown that the music-induced increase in aerobic performance time may be mainly attributed to the energizing and encouraging effects of music. It is said that listening to music during exercise may narrow down the attention and distract the individual from the exercise-induced feeling of fatigue, because the data processing capacities are limited and this changes the individual's RPE [29, 31].

Combining exercise with music may increase cognitive provocations via generating motivation [32]. Also, music may replace the sensory data pertaining to physical activities transmitted to the central nervous system, increase the efficiency of performance and improve the activity-induced agitation [33]. Research has shown that muscular tension changes as the type of music changes: exciting music increases but relaxing music decreases the muscular tension [34]. Relaxing music may reduce muscular activity during exercise, but exciting music increases the pedaling and running time and delays fatigue; therefore, it encourages the individual to do the exercise and allows him/her to lengthen the exercise time [9].

In conclusion, the results of the present study showed that the mean fatigue onset time and mean heart rate were significantly higher in the participants while they were listening to exciting and arbitrary music compared to relaxing and no music. RPE was significantly lower in the participants while they were listening to exciting and arbitrary music, compared with relaxing and no music. However, listening to different types of music did not significantly affect systolic and diastolic blood pressure.

## References

- 1-kendelhardt AR (2003). Effect of live music on exercise duration, negative verbalization, and self-perception of pain, anxiety, and rehabilitation level of physical therapy patients. Electronic Theses, Treatises and Dissertations, PP: 31-85. <http://diginole.lib.fsu.edu/etd/3185>.
- 2-Karageorghis, CI (1999). Music in sport and exercise: Theory and practice. *The Sport Journal* 2(2). Retrieved March 28, 2007, from <http://www.thesportjournal.org/1999Journal/Vol2-No2/Music.asp>.
- 3-Szmedra L, Bacharach DW (1998). Effect of music on rate of perceived exertion, plasma lactate, norepinephrine and cardiovascular Homodynamic during Treadmill running. *Int J sports med* 19: 32-37.
- 4-Karageorghis CL, Drew KM, Terry PC (1996). Effect of pretest stimulative and sedative music on grip strength. perceptual and motor skill 83: 1347-52.
- 5-Khorram R (2005). Investigating the effects of listening to exciting and classical music on running time, salivary cortisol concentration and RPE during one session training to the exhaustion limit in the male students of physical education. Unpublished MA thesis, Tehran-center branch, Islamic Azad University, Iran, PP: 26-45.
- 6-Boutcher SH, Trenske M (1990). The effects of sensory

- deprivation and music on perceived exertion and affect during exercise. *J Sport Exe Psycho* 2: 169-176.
- 7-Nethery VM (2002). Competition between internal and external sources of information during exercise: influence on RPE and the impact of the exercise load. *J Sport Med and Physical Fitness* 42(2): 172-8.
  - 8-Costasi K, Terry PC (1997). The psychophysical effects of music in sport and exercise: A review. *Journal of Sport Behavior* 20:54-68.
  - 9-Copland BL, Franks BD (1991). Effects of type and intensities of background music on treadmill endurance. *Journal of sport medicine and physical fitness* 51(1): 100-103.
  - 10-Brenda L, Copland B, Don F (1991). Effect of types and intensities of background music on treadmill endurance. *J Sport Med phys Fit* 31 (1): 100-103.
  - 11-Fathollahi Z. (2006). The effects of listening to two types of music on running time, salivary cortisol and RPE during one session training in active women. Unpublished MA thesis, Tehran University, Iran, pp: 26-47.
  - 12-Yamashita S, Iwai K, Akimoto T, Sugawara J, Kono I (2006). Effect of music during exercise on RPE, heart rate and the autonomic nervous system. *J sport med phys fitness* 46(3): 423-430.
  - 13-MohammedZadeh H, Ahmedi A (2008). Effects of rapid music on exercise performance, and perceived pressure and increasing activity to exhaustion in untrained. *J Movement* 38: 147-159.
  - 14-Torabi F (2009). The effects of external intervention (music) on some fatigue indicators in healthy young women. *Journal of sport science research* 22: 51-62.
  - 15-Liptak V (1979). Psycho-physiological effects of pop music on cardiovascular parameters at rest during exertion. *Wien. Med Wochenschr* 10: 268-275.
  - 16-Davis WB, Thaut MH (1989). The influence of proffered relaxation music on measure of state anxiety, relaxation and physiological response. *J music therapy* 26:168-187.
  - 17-Brownley KA, Murray RG, Hackney AC (1995). Effect of music on physiological and affective responses to graded treadmill exercise in trained and untrained runner. *Inter J Psychophysiology* 19(3): 193-201.
  - 18-Knight WE, Richard, NS (2001). Relaxing music prevents stress-induced increases in subjective anxiety systolic blood pressure and heart rate in healthy male and females. *J music Therapy* 33(4): 254-272.
  - 19-Nikbakhsh R (1996). The effects of sensory deprivation and music on RPE and temperament during training. Unpublished MA thesis, Islamic Azad University, Tehran center branch, Iran, pp: 21-51.
  - 20-Szabo A, Small A, Leigh A (1999). The effects of slow and fast rhythm classical music on progressive cycling to voluntary physical exhaustion". *Journal of sports Med and Physical fitness* 39: 220-225.
  - 21-Crust L (2004). Affect of familiar and unfamiliar a synchronous music on treadmill walking endurance. perceptual and motor skill 99: 361-368.
  - 22-Yamamoto T, Ohkuwa T, Itoh H, Kitoh M, Terasaw J, Tsuda T, Kitagawa S, Sato Y (2003). Effects of pre-exercise listening to slow and fast rhythm music on supra maximal cycle performance and selected metabolic variables. *Arch Physiol Biochem* 111(3): 211-4.
  - 23-Sears W (1975). The effect of music on muscle tonus. In E.G.Gaston. (Ed), *music therapy* (pp. 199-205). Lawrence, KS: Allen Press.
  - 24-Carla H, Rachel K (1996). Effect of music on cardiovascular performance during on treadmill endurance. *Journal of music and imagery of physical Fitness* 31: 100-103.
  - 25-Lorch CA, Lorch V, Diefendorf A, Earl PW (1994). Effect of simulative and sedative music on systolic blood pressure, heart rate, and respiratory rate in premature infants. *Journal of music therapy* 31(2): 105-118.
  - 26-Ghaderi M Aghaalinezhad H, Azarbayjani MA (2008). Investigating the effects of exciting and relaxing music on aerobic performance, RPE and salivary cortisol concentration in male student athletes. *Olympic* 41: 17-23.
  - 27-Ahmedi A (2007). The effects of techno music on RPE and performance in cumulative activity to the exhaustion limit in both the individuals who have and those who have not taken exercise. Unpublished MA thesis, Tehran University, Iran, pp: 31-44.
  - 28-Janathan D, Meeks RT (2002). Music enhances performance but not physiological recovery following exercise, (paper accepted for presentation at a veal Meeting of the American psychological society. New Orleans).
  - 29-Maraki M, Tsoliou F, Pitsitadis YP, Malkova D, Mutrie N, Higgins S (2005). Acute effects of single exercise class appetite, energy intake and mood. Is there time of day effect? *Appetite* 9; [E pub ahead of print].
  - 30-Rejeski WJ (1985). perceived exertion: an active or passive process? *J. Sport Psychology* 7: 371-378.
  - 31-Wales DN (1986). The effects of tempo and disposition in music on perceived exertion, brain waves and mood during aerobic exercise. Master's thesis, Pennsylvania state university, pp:23-38.
  - 32-Emery CF (2004). Fitness: A little music with exercise boosts brain power. *Study Suggests* [1], *Pain & Central nervous system week*, P: 30.
  - 33-Hayakawa YH, Miki K, Akada KT (2000). Effect of music on mood during bench stepping exercise. *percept. Mot Skills* 1: 307-314.
  - 34-Mark IJ, Alpert I, Elliot MN (2005). Purchase occasion influence on the role of music in advertising. *J Business Research* 58: 369-376.